



Mitchell E. Daniels, Jr.
Governor

Judith A. Monroe, M.D.
State Health Commissioner

Indiana State Department of Health

An Equal Opportunity Employer

DATE: March 24, 2008

TO: All Local Health Departments
Attn: Chief Food Inspection Officer

FROM: A. Scott Gilliam, MBA, CFSP
Manager, Food Protection Program

SUBJECT: FDA Warns of Salmonella Risk with Cantaloupes from Agropecuaria Montelibano

From the information provided by FDA the import alert product, cantaloupes from Agropecuaria Montelibano, a Honduran company, were not distributed in the State of Indiana. The cantaloupes were distributed to neighboring States of Illinois and Ohio. In addition, if any cantaloupes are found from this grower, please notify Dan Gala at this office at 317-233-7360.

Advisory; Detain cantaloupes; no notification of affected product required.

FDA NEWS RELEASE

Media Inquiries: Stephanie Kwisnek, 301-827-0955, stephanie.kwisnek@fda.hhs.gov
Consumer Inquiries: 888-INFO-FDA

FDA Warns of Salmonella Risk with Cantaloupes from Agropecuaria Montelibano *The agency detains products from the Honduran manufacturer*

The U.S. Food and Drug Administration has issued an import alert regarding entry of cantaloupe from Agropecuaria Montelibano, a Honduran grower and packer, because, based on current information, fruit from this company appears to be associated with a *Salmonella* Litchfield outbreak in the United States and Canada. The import alert advises FDA field offices that all cantaloupes shipped to the United States by this company are to be detained.

In addition, the FDA has contacted importers about this action and is advising U.S. grocers, food service operators, and produce processors to remove from their stock any cantaloupes from this company. The FDA also advises consumers who have recently bought cantaloupes to check with the place of purchase to determine if the fruit came from this specific grower and packer. If so, consumers should throw away the cantaloupes.

To date, the FDA has received reports of 50 illnesses in 16 states and nine illnesses in Canada linked to the consumption of cantaloupes. No deaths have been reported; however, 14 people have been hospitalized. The states are Arizona, California, Colorado, Georgia, Illinois, Missouri, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Tennessee, Utah, Washington, and Wisconsin.

Epidemiology Resource Center
2525 N. Shadeland Ave. Suite E3, Indianapolis, IN 46219
317.356.7190 ext. 253

Laboratories
635 North Barhill Dr. Room 2031, Indianapolis, IN 46202
317.233.8000

Weights & Measures
2525 N. Shadeland Ave. Suite D3, Indianapolis, IN 46219
317.356.7078 ext. 221

The FDA is taking this preventive measure while the agency continues to investigate this outbreak in cooperation with the Centers for Disease Control and Prevention and state partners. Such intervention is a key component of FDA's [Food Protection Plan](#).

Symptoms of foodborne Salmonella infection include nausea, vomiting, fever, diarrhea, and abdominal cramps. In persons with poor health or weakened immune systems, Salmonella can invade the bloodstream and cause life-threatening infections. Individuals who have recently eaten cantaloupe and experienced any of these symptoms should contact their health care professional.

The FDA recommends that consumers take the following steps to reduce the risk of contracting Salmonella or other foodborne illnesses from cantaloupes:

- Purchase cantaloupes that are not bruised or damaged. If buying fresh-cut cantaloupe, be sure it is refrigerated or surrounded by ice.
- After purchase, refrigerate cantaloupes promptly.
- Wash hands with hot, soapy water before and after handling fresh cantaloupes.
- Scrub whole cantaloupes by using a clean produce brush and cool tap water immediately before eating. Don't use soap or detergents.
- Use clean cutting surfaces and utensils when cutting cantaloupes. Wash cutting boards, countertops, dishes, and utensils with hot water and soap between the preparation of raw meat, poultry, or seafood and the preparation of cantaloupe.
- If there happens to be a bruised or damaged area on a cantaloupe, cut away those parts before eating it.
- Leftover cut cantaloupe should be discarded if left at room temperature for more than two hours.
- Use a cooler with ice or use ice gel packs when transporting or storing cantaloupes outdoors.